

Superconducting cosmic strings as Gamma Ray Burst engines

V. Berezhinsky¹⁾, B. Hnatyk^{1),2)} and A. Vilenkin³⁾

¹⁾*INFN, Laboratori Nazionali del Gran Sasso, I-67010 Assergi (AQ), Italy*

²⁾*Institute for Applied Problems in Mechanics and Mathematics, NASU, Naukova 3b, Lviv-53, 290053, Ukraine*

³⁾*Physics Department, Tufts University, Medford, MA 02155, USA*

Cusps of superconducting strings can serve as GRB engines. A powerful beamed pulse of electromagnetic radiation from a cusp produces a jet of accelerated particles, whose propagation is terminated by the shock responsible for GRB. A single free parameter, the string scale of symmetry breaking $\eta \sim 10^{14}$ GeV, together with reasonable assumptions about the magnitude of cosmic magnetic fields and the fraction of volume that they occupy, explains the GRB rate, duration and fluence, as well as the observed ranges of these quantities. The wiggles on the string can drive the short-time structures of GRB. This model predicts that GRBs are accompanied by strong bursts of gravitational radiation which should be detectable by LIGO, VIRGO and LISA detectors.

Models of gamma ray bursts (GRBs) face the problem of explaining the tremendous energy released by the central engine [1]. In the case of isotropic emission, the total energy output should be as high as 4×10^{54} ergs. Strongly beamed emission is needed for all known engine models, such as mergers and hypernovae, but such extreme beaming is difficult to arrange. In this paper we show that emission of pulsed electromagnetic radiation from cusps of superconducting cosmic strings naturally solves this problem and explains the observational GRB data using only one engine parameter.

Cosmic strings are linear defects that could be formed at a symmetry breaking phase transition in the early universe [2]. Strings predicted in most grand unified models respond to external electromagnetic fields as thin superconducting wires [3]. As they move through cosmic magnetic fields, such strings develop electric currents. Oscillating loops of superconducting string emit short bursts of highly beamed electromagnetic radiation and high-energy particles [4,5].

The idea that GRBs could be produced at cusps of superconducting strings was first suggested by Babul, Paczynski and Spergel [6] (BPS) and further explored by Paczynski [7]. They assumed that the bursts originate at very high redshifts ($z \sim 100 - 1000$), with GRB photons produced either directly or in electromagnetic cascades developing due to interaction with the microwave background. This model requires the existence of a strong primordial magnetic field to generate the string currents.

As it stands, the BPS model does not agree with observations. The observed GRB redshifts are in the range $z \lesssim 3$, and the observed duration of the bursts ($10^{-2}s \lesssim \tau \lesssim 10^3s$) is significantly longer than that predicted by the model. On the theoretical side, our understanding of cosmic string evolution and of the GRB generation in relativistic jets have considerably evolved since the BPS papers were written. Our goal in this paper is to revive the BPS idea taking stock of these recent advances.

As in the BPS model we shall use the cusp of a superconducting string as the central engine in GRB. It provides the tremendous engine energy naturally beamed. Our main observation is that putting superconducting cusps in a different environment, the magnetized plasma at a relatively small redshift z , results in a different mechanism of gamma radiation, which leads to a good agreement with GRB observational data.

GRB radiation in our model arises as follows. Low-frequency electromagnetic radiation from a cusp loses its energy by accelerating particles of the plasma to very large Lorentz factors. Like the initial electromagnetic pulse, the particles are beamed and give rise to a hydrodynamical flow in the surrounding gas, terminated by a shock, as in the standard fireball theory of GRB [8] (for a review see [1]). We shall assume that cosmic magnetic fields were generated at moderate redshifts (*e.g.*, in young galaxies during the bright phase of their evolution [9]). The string symmetry breaking scale η will be the only string parameter used in our calculations. With reasonable assumptions about the magnitude of cosmic magnetic fields and the fraction of volume in the Universe that they occupy, this parameter is sufficient to account for all main GRB observational quantities: the duration τ_{GRB} , the rate of events $\dot{N}_{GRB}$, and the fluence S .

We begin with the description of some properties of strings, which will be used below (for a review see [2]).

A horizon-size volume at any time t contains a few long strings stretching across the volume and a large number of small closed loops. The typical length of a loop at cosmological time t and the loop number density are given by

$$l \sim \alpha t, \quad n_l(t) \sim \alpha^{-1} t^{-3} \quad (1)$$

The exact value of the parameter α in (1) is not known. We shall assume, following [10], that α is determined by the gravitational backreaction, so that $\alpha \sim k_g G \mu$, where $k_g \sim 50$ is a numerical coefficient, G is Newton's

constant, $\mu \sim \eta^2$ is the mass per unit length of string, and η is the symmetry breaking scale of strings.

The loops oscillate and lose their energy, mostly by gravitational radiation. For a loop of invariant length l [11], the oscillation period is $T_l = l/2$ and the lifetime is $\tau_l \sim l/k_g G\mu$.

An electric field E applied along a superconducting string generates an electric current. A superconducting loop of string oscillating in a magnetic field B acts as an *ac* generator and develops an *ac* current of amplitude

$$J_0 \sim e^2 B l. \quad (2)$$

The local value of the current in the loop can be greatly enhanced in near-cusp regions where, for a short period of time, the string reaches a speed very close to the speed of light. Cusps tend to be formed a few times during each oscillation period. Near a cusp, the string gets contracted by a large factor, its rest energy being turned into kinetic energy. The density of charge carriers, and thus the current, are enhanced by the same factor. The contraction factor increases as one approaches the point of the cusp.

The growth of electric current near the cusp due to string contraction is terminated at a critical value J_{max} when the energy of charge carriers becomes comparable to that of the string itself, $(J/e)^2 \sim \mu$. This gives

$$J_{max} \sim e\eta, \quad \gamma_{max} \sim (e\eta/J_0). \quad (3)$$

Alternatively, the cusp development can be terminated by small-scale wiggles on the string [12]. If the wiggles contribute a fraction $\epsilon \ll 1$ to the total energy of the string, then the maximum Lorentz factor is less than (3), and is given by $\gamma_{max} \sim \epsilon^{-1/2}$. The actual value of γ_{max} is not important for most of the following discussion.

Due to the large current, the cusp produces a powerful pulse of electromagnetic radiation. The total energy of the pulse is given by [4,5] $\mathcal{E}_{em}^{tot} \sim 2k_{em}J_0J_{max}l$, where $l \sim \alpha t$ is the length of the loop, and the coefficient $k_{em} \sim 10$ is taken from numerical calculations [4]. This radiation is emitted within a very narrow cone of opening angle $\theta_{min} \sim 1/\gamma_{max}$ (for relativistic beaming in GRB see [13]). The angular distribution of radiated energy at larger angles is given by [4]

$$d\mathcal{E}_{em}/d\Omega \sim k_{em}J_0^2l/\theta^3, \quad (4)$$

We shall adopt the following simple model of cosmic magnetic fields. We shall assume that magnetic fields were generated at some $z \sim z_B$ and then remained frozen in the extragalactic plasma, with

$$B(z) = B_0(1+z)^2, \quad (5)$$

where B_0 is the characteristic field strength at the present time. For numerical estimates below we shall use $z_B \sim 4$, $B_0 \sim 10^{-7}$ G, and assume that the fraction of volume of

the universe occupied by magnetized plasma is $f_B \sim 0.1$. We shall also assume that the universe is spatially flat and is dominated by non-relativistic matter.

We shall now estimate the physical quantities characterizing GRBs powered by superconducting strings. For a GRB originating at redshift z and seen at angle θ with respect to the string velocity at the cusp, we have from Eqs.(2)-(5)

$$d\mathcal{E}_{em}/d\Omega \sim k_{em}e^4\alpha^3t_0^3B_0^2(1+z)^{-1/2}\theta^{-3}, \quad (6)$$

where t_0 is the present age of the Universe. The Lorentz factor of the relevant string segment near the cusp is $\gamma \sim 1/\theta$. The duration of the cusp event as seen by a distant observer is [6]

$$\tau_c \sim (1+z)(\alpha t/2)\gamma^{-3} \sim (\alpha t_0/2)(1+z)^{-1/2}\theta^3. \quad (7)$$

One can expect that the observed duration of GRB is $\tau_{GRB} \sim \tau_c$. This expectation will be justified by the hydrodynamical analysis below.

The fluence, defined as the total energy per unit area of the detector, is [7]

$$S \sim (1+z)(d\mathcal{E}_{em}/d\Omega)d_L^2(z), \quad (8)$$

where $d_L(z) = 3t_0(1+z)^{1/2}[(1+z)^{1/2}-1]$ is the luminosity distance.

The rate of GRBs originating at cusps in the redshift interval dz and seen at an angle θ in the interval $d\theta$ is given by

$$d\dot{N}_{GRB} \sim f_B \cdot \frac{1}{2}\theta d\theta(1+z)^{-1}\nu(z)dV(z). \quad (9)$$

Here, $\nu(t) \sim n_l(t)/T_l \sim 2\alpha^{-2}t^{-4}$ is the number of cusp events per unit spacetime volume, $T_l \sim \alpha t/2$ is the oscillation period of a loop, $dV = 54\pi t_0^3[(1+z)^{1/2}-1]^2(1+z)^{-11/2}dz$ is the proper volume between redshifts z and $z+dz$, and we have used the relation $dt_0 = (1+z)dt$.

Since different cusp events originate at different redshifts and are seen at different angles, our model automatically gives a distribution of durations and fluences of GRBs. The angle θ is related to the Lorentz factor of the relevant portion of the string as $\theta \sim 1/\gamma$, and from Eqs.(6),(8) we have

$$\gamma(z; S) \sim \gamma_0\alpha_{-8}^{-1}S_{-8}^{1/3}B_{-7}^{-2/3}[(\sqrt{1+z}-1)^2\sqrt{1+z}]^{1/3}. \quad (10)$$

Here, $\gamma_0 \approx 190$, $\alpha_{-8} = \alpha/10^{-8}$, and the fluence S and the magnetic field B_0 are expressed as $S = S_{-8} \cdot 10^{-8}$ erg/cm² and $B = B_{-7} \cdot 10^{-7}$ G.

Very large values of $\gamma \sim \gamma_{max}$, which correspond (for a given redshift) to largest fluences, may not be seen at all because the radiation is emitted into a too narrow solid angle and the observed rates of these events

are too small. The minimum value $\gamma(z; S_{min})$ is determined by the smallest fluence that is observed, $S_{min} \sim 2 \cdot 10^{-8} \text{ erg/cm}^2$. Another limit on γ , which dominates at small z , follows from the condition of compactness [1] and is given by $\gamma \gtrsim 100$ (see below).

The total rate of GRBs with fluence larger than S is obtained by integrating Eq.(9) over θ from $\gamma_{max}^{-1}(z)$ to $\gamma^{-1}(z; S)$ and over z from 0 to $\min[z_m; z_B]$, with z_m from $\gamma_{max}(z_m) = \gamma(z_m; S)$. For relatively small fluences, $S_{-8} < S_c = 0.03(\gamma_{max}(0)\alpha_{-8}/\gamma_0)^3 B_{-7}^2$, $z_B < z_m$ and we obtain

$$\begin{aligned} \dot{N}_{GRB}(> S) &\sim \frac{f_B}{2\alpha^2 t_0^4} \int_0^{z_B} dV(z)(1+z)^5 \gamma^{-2}(z; S) \\ &\sim 3 \cdot 10^2 S_{-8}^{-2/3} B_{-7}^{4/3} \text{ yr}^{-1}. \end{aligned} \quad (11)$$

Remarkably, this rate in our model does not depend on any string parameters and is determined (for a given value of S) almost entirely by the magnetic field B_0 . The predicted slope $\dot{N}_{GRB}(> S) \propto S^{-2/3}$ is in a reasonable agreement with the observed one $\dot{N}_{obs}(> S) \propto S^{-0.55}$ at relatively small fluences [14].

For large fluences $S_{-8} > S_c$, integration of Eq.(9) gives $\dot{N}_{GRB}(> S) \propto S^{-3/2}$. Observationally, the transition to this regime occurs at $S_{-8} \sim 10^2 - 10^3$. This can be accounted for if the cusp development is terminated by small-scale wiggles with fractional energy in the wiggles $\epsilon \sim 10^{-7} \alpha_{-8}^2 B_{-7}^{4/3}$. Alternatively, if γ_{max} is determined by the back-reaction of the charge carriers, Eq.(3), then the regime (11) holds for larger S_{-8} , and observed steepening of the distribution at large S can be due to the reduced efficiency of BATSE to detection of bursts with large γ . Indeed, large γ results in a large Lorentz factor γ_{CD} of the emitting region (see below), and at $\gamma_{CD} \gtrsim 10^3$ photons start to escape from the BATSE range.

The duration of a GRBs originating at redshift z and having fluence S is readily calculated as

$$\tau_{GRB} \approx 200 \frac{\alpha_{-8}^4 B_{-7}^2}{S_{-8}} (1+z)^{-1} (\sqrt{1+z} - 1)^{-2} s \quad (12)$$

Estimated from Eq.(7), $\tau_{GRB}^{max} \sim 10^3 \alpha_{-8} s$, while from Eq.(12) using $S_{max} \sim 1 \cdot 10^{-4} \text{ erg/cm}^2$ and $z \sim z_B \sim 4$, one obtains $\tau_{GRB}^{min} \sim 3 \alpha_{-8}^4 B_{-7}^2 \text{ ms}$. This range of τ_{GRB} agrees with observations for $\alpha_{-8} \sim 1$ (*i.e.* $\eta \sim 2 \cdot 10^{14} \text{ GeV}$) and $B_{-7} \sim 1$.

Let us now turn to the hydrodynamical phenomena in which the gamma radiation of the burst is actually generated. The low-frequency electromagnetic pulse interacting with surrounding gas produces an ultrarelativistic beam of accelerated particles. This is the dominant channel of energy loss by the pulse. The beam of high energy particles pushes the gas with the frozen magnetic field ahead of it, producing an external shock in surrounding plasma and a reverse shock in the beam material, as in the case of ‘‘ordinary’’ fireball (for a review see [1]).

The difference is that the beam propagates with a very large Lorentz factor $\gamma_b \gg \gamma$ where γ is the Lorentz factor of the cusp. (The precise value of γ_b is not important for this discussion; it will be estimated in a subsequent publication [15].) Another difference is that the beam propagates in a very low-density gas. The beam can be regarded as a narrow shell of relativistic particles of width $\Delta \sim l/2\gamma^3$ in the observer’s frame.

The gamma radiation of the burst is produced as synchrotron radiation of electrons accelerated by external and reverse shocks. Naively, the duration of synchrotron radiation, *i.e.* τ_{GRB} , is determined by the thickness of the shell as $\tau_{GRB} \sim \Delta$. This is confirmed by a more detailed analysis, as follows. The reverse shock in our case is ultrarelativistic [16,1]. The necessary condition for that, $\rho_b/\rho < \gamma_b^2$, is satisfied with a wide margin (here ρ_b is the baryon density in the beam and ρ is the density of unperturbed gas). In this case, the shock dynamics and the GRB duration are determined by two hydrodynamical parameters [1]. They are the thickness of the shell Δ and the Sedov length, defined as the distance travelled by the shell when the mass of the snow-ploughed gas becomes comparable to the initial energy of the beam. The latter is given by $l_{Sed} \sim (\mathcal{E}_{iso}/\rho)^{1/3}$.

The reverse shock enters the shell and, as it propagates there, it strongly decelerates the shell. The synchrotron radiation occurs mainly in the shocked regions of the shell and of the external plasma. The surface separating these two regions, the contact discontinuity (CD) surface, propagates with the same velocity as the shocked plasma, where the GRB radiation is produced.

The synchrotron radiation ceases when the reverse shock reaches the inner boundary of the shell. This occurs at a distance $R_\Delta \sim l_{Sed}^{3/4} \Delta^{1/4}$ when the Lorentz factor of the CD surface is $\gamma_{CD} \sim (l_{Sed}/\Delta)^{3/8} \sim 0.1 B_{-7}^{1/4} n_{-5}^{-1/8} (1+z)^{1/2} \gamma^{3/2}$, where n_{-5} is the ambient baryon number density in units of 10^{-5} cm^{-3} . Note that these values do not depend on the Lorentz factor of the beam γ_b and are determined by the cusp Lorentz factor γ . The size of the synchrotron emitting region is of the order R_Δ , and the Lorentz factor of this region is equal to γ_{CD} . The compactness condition [1] requires $\gamma_{CD} \gtrsim 10^2$, which yields $\gamma \gtrsim 10^2$ used above. The duration of GRB is given by

$$\tau_{GRB} \sim R_\Delta / 2\gamma_{CD}^2 \sim l/2\gamma^3, \quad (13)$$

i.e. it is equal to the duration of the cusp event given by Eq.(7). The energy that goes into synchrotron radiation is comparable to the energy of the electromagnetic pulse.

We conclude with several remarks on our GRB model.

(i) Magnetic fields in our scenario could be generated in young galaxies during the bright phase of their evolution [9] and then dispersed by galactic winds in the intergalactic space. One expects that at present the fields are concentrated in the filaments and sheets of the large-

scale structure [17,18]. With sheets of characteristic size $L \sim (20 - 50)h^{-1}$ Mpc and thickness $D \sim 5h^{-1}$ Mpc, we have $f_B \sim D/L \sim 0.1$. The average field strength can be estimated from the equipartition condition as $B_0 \sim 10^{-7}$ G [17]. We expect these values to be accurate within an order of magnitude.

(ii) We emphasize that our model involves a number of simplifying assumptions. All loops at cosmic time t were assumed to have the same length $l \sim \alpha t$, while in reality there should be a distribution $n(l, t)$. The evolution law (5) for $B(z)$ and the assumption of $f_B = \text{const}$ are also oversimplified. A more realistic model should also account for a spatial variation of B . As a result, the correlation between τ and S suggested by Eq.(12) will tend to be washed out. Finally, we disregarded the possibility of a nonzero cosmological constant which would introduce some quantitative changes in our estimates.

(iii) Apart from the reverse shock instability, small-scale wiggles on strings can naturally produce short-time variation in GRBs. These wiggles, acting like mini-cusps, produce a sequence of successive fireballs, which are the usual explanation of GRB short-time structure [1].

(iv) Cusps reappear on a loop, producing nearly identical GRBs with a period $T_l \sim \alpha t/2 \sim 50(1+z)^{-1/2}$ yr. In a more realistic model, some fraction of loops would have lengths smaller than αt and shorter recurrence periods.

(v) Our model predicts that GRBs should be accompanied by strong bursts of gravitational radiation. The angular distribution of the gravitational wave energy around the direction of the cusp is [19] $d\mathcal{E}_g/d\Omega \sim G\mu^2/\theta$, and the dimensionless amplitude of a burst of duration τ originating at redshift z can be estimated as

$$h \sim k_g(G\mu)^2(\tau/\alpha t_0)^{1/3}(1+z)^{-1/3}[(1+z)^{1/2} - 1]^{-1}, \quad (14)$$

or $h \sim 10^{-21} \alpha_{-8}^{5/3} z^{-1} (\tau/1s)^{1/3}$ for $z \lesssim 1$. Here, we have used the relation $F_g \sim h^2/G\tau^2 \sim (1+z)(d\mathcal{E}_g/d\Omega)d_L^{-2}(z)$ for the gravitational wave flux and Eq.(7) for the burst duration τ . These gravitational wave bursts are much stronger than expected from more conventional sources and should be detectable by the planned LIGO, VIRGO and LISA detectors.

(vi) GRBs have been suggested as possible sources of the observed ultrahigh-energy cosmic rays (UHECR) [20,21]. This idea encounters two difficulties. If GRBs are distributed uniformly in the universe, UHECR have a classical Greisen-Zatsepin-Kuzmin (GZK) cutoff, absent in the observations. The acceleration by an ultra-relativistic shock is possible only in the one-loop regime (*i.e.* due to a single reflection from the shock) [22]. For a standard GRB with a Lorentz factor $\gamma_{sh} \sim 300$ it results in the maximum energy $E_{max} \sim \gamma_{sh}^2 m_p \sim 10^{14}$ eV, far too low for UHECR.

Our model can resolve both of these difficulties, assuming that γ_{max} is determined by the current back-

reaction, Eq.(3).

If the magnetic field in the Local Supercluster (LS) is considerably stronger than outside, then the cusps in LS are more powerful and the GZK cutoff is less pronounced.

Near-cusp segments with large Lorentz factors produce hydrodynamical flows with large Lorentz factors, *e.g.* $\gamma \sim 2 \cdot 10^4$ corresponds to $\gamma_{CD} \sim 3 \cdot 10^5$ and $E_{max} \sim \gamma_{CD}^2 m_p \sim 1 \cdot 10^{20}$ eV. Protons with such energies are deflected in the magnetic field of LS and can be observed, while protons with much higher energies caused by near-cusp segments with $\gamma \gtrsim 10^5$ are propagating rectilinearly and generally are not seen. Further details of this scenario will be given elsewhere [15].

We are grateful to Ken Olum for useful discussions. The work of AV was supported in part by the National Science Foundation and the work of VB and BH by INTAS through grant No 1065.

-
- [1] T. Piran astro-ph/9810256
 - [2] A. Vilenkin and E.P.S. Shellard, *Cosmic strings and other topological defects*, Cambridge University Press, Cambridge, 1994.
 - [3] E. Witten, Nucl. Phys. **B249**, 557 (1985).
 - [4] A. Vilenkin and T. Vachaspati, Phys. Rev. Lett. **58**, 1041 (1987).
 - [5] D.N. Spergel, T. Piran and J. Goodman, Nucl. Phys. **B291**, 847 (1987).
 - [6] A. Babul, B. Paczynski and D.N. Spergel, Ap. J. Lett. **316**, L49 (1987).
 - [7] B. Paczynski, Ap. J. **335**, 525 (1988).
 - [8] P. Meszaros and M.J. Rees, MNRAS, **258**, 41 (1992).
 - [9] R.B. Partridge and P.J. Peebles, Ap.J., **147**, 868 (1967).
 - [10] D. Bennett and F. Bouchet, Phys. Rev. Lett, **60**, 257 (1988).
 - [11] The actual length of the loop changes as the loop oscillates. The invariant length is defined as $l = E/\mu$, where E is the loop's energy in its center-of-mass frame.
 - [12] B. Carter, Phys. Rev. **D41**, 3869 (1990); A. Vilenkin, Phys. Rev. **D41**, 3038 (1990).
 - [13] S. Mao and I. Yi, Ap.J. **424**, L131 (1994).
 - [14] Catalog BATSE 4B, cited in G. Pugliese et al astro-ph/9912019.
 - [15] V. Berezhinsky, B. Hnatyk and A. Vilenkin, in preparation.
 - [16] S. Kobayashi, T. Piran and R. Sari, astro-ph/9803217
 - [17] D. Ryu, H. Kang and P.L. Biermann, astro-ph/9803275.
 - [18] G.R. Farrar and T. Piran, astro-ph/9906431.
 - [19] T. Vachaspati and A. Vilenkin, Phys. Rev. **D31**, 3052 (1985).
 - [20] M. Vietri, Ap.J. **453**, 883 (1995).
 - [21] E. Waxman, Phys. Rev. Lett. **75**, 386 (1995).
 - [22] Y.A. Gallant and A. Achterberg, MNRAS, **305**, L6 (1999)